

MULTIVA® Active Cat

Multi-Vitamin-Mineral Supplement with Amino Acids, Omega-3, Omega-6 and Digestive Enzymes Formulated into Highly Palatable Chews.



Data Sheet

Active Ingredients (per Chew):

Taurine	200 mg	Vitamin B ₅ (Pantothenic Acid)	5 mg
Methionine	10 mg	Vitamin B ₆ (Pyridoxine)	2 mg
Cystine	5 mg	Vitamin B ₇ (Biotin)	2 mcg
DHA (Omega-3)	5.5 mg	Vitamin B ₉ (Folic Acid)	2 mcg
EPA (Omega-3)	8.2 mg	Vitamin B ₁₂ (Cobalamin)	2 mcg
Linoleic Acid (Omega-6)	25 mg	Coline (before Vitamina B _p)	5 mg
Calcium	50 mg	Inositol (before Vitamina B _m)	2 mg
Phosphorus	25 mg	PABA (before Vitamin B ₁₀)	2 mg
Potassium	0.25 mg	Vitamin C (Ascorbic Acid)	20 mg
Magnesium	1.5 mg	Vitamin D ₃	50 IU
Iron	1.5 mg	Vitamin E	5 IU
Copper	30 mcg	Betaine	10 mg
Zinc	1 mg	Lecithin	5 mg
Iodine	40 mcg	Bromelain	0.38 UGD
Selenium	3.5 mcg	Lipase	1.08 UUSP
Vitamin A	900 IU	Protease	13.6 UUSP
Vitamin B ₁ (Thiamine)	2 mg	Pepsin	2.55 UNF
Vitamin B ₂ (Riboflavin)	2 mg	Papain	1.7 UUSP
Vitamin B ₃ (Niacin)	5 mg		

Composition (in descending order): chicken flavour, yeast, glycerol, taurine, calcium phosphate, soy lecithin, whey, sweetener (maltodextrin, sodium alginate, calcium sulphate), rapeseed oil, L-ascorbyl-2-polyphosphate, calcium carbonate, safflower oil, fish oil, choline chloride, betaine hydrochloride, methionine, vitamin E, iron proteinate, zinc proteinate, calcium pantothenate, cystine, niacinamide, bromelain, magnesium oxide, riboflavin, pyridoxine hydrochloride, monohydrate thiamine, PABA, inositol, vitamin B₁₂, vitamin A (acetate), propionic acid, mixed tocopherols, potassium chloride, pancreatin, copper proteinate, sodium selenite, vitamin D₃, biotin, calcium iodate, folic acid, pepsin, papain.

Additives (per kg):

Vitamins: E672 vitamin A (acetate) 552 mg; E671 vitamin D₃ 48mg.
Oligoelements: E1 iron (proteinate) 600 ppm; E2 iodine (calcium iodate) 16 ppm; E4 copper (proteinate) 12 ppm mg; E6 zinc (proteinate) 400 ppm; E8 selenium (sodium selenite) 1.4 ppm.

Analytical Constituents: Crude protein 26.6%; fat content 13.05%; crude fibre 1.58%; crude ash 14.50%; moisture content 5.94%.

Mechanism of Action:

MULTIVA® Active Cat contains a unique combination of vitamins, minerals, amino acids, Omega-3, Omega-6 and digestive enzymes that act synergistically to ensure the macro and micronutrients necessary for the optimal functioning of the organism.

- The intake of vitamins is essential for proper functioning of organs and systems, however it is important that the proportions are adequate and balanced.
 - **Vitamin A** contributes to the normal metabolism of iron, maintenance of natural barriers such as skin



Features

Complete formula of 37 essential nutrients: Vitamins, Minerals, Omega-3 and Omega-6, Taurine, Cystine, Methionine and Digestive Enzymes.

Contains macro and micronutrients that act synergistically for optimal functioning of the organism.

Combats nutritional deficiencies associated with insufficient intake or increased body demands.

Increases the appetite and activity of animals that are listless, apathetic or reluctant to eat.

Nutritional support in stressed, sick or recuperating animals.

Helps to strengthen the immune system, improving defences in situations of stress, convalescence...

Recommended in animals with high nutritional needs.

Boosts the health of geriatric animals.



MULTIVA® Active Cat

Multi-Vitamin-Mineral Supplement with Amino Acids, Omega-3, Omega-6 and Digestive Enzymes Formulated into Highly Palatable Chews.



Data Sheet

Prevents and combats excessive hair loss, other dermatological problems and some intestinal pathologies such as occur with poor digestion and weight loss.

Recommended by veterinarians
–Complete, balanced and effective.

Highly palatable Chews – Facilitate daily intake.

Suitable for all breeds and ages.

- and mucosa, normal functioning of the immune system and maintenance of healthy vision
- **Group B Vitamins** have neuroprotective, antioxidant properties and are essential for the synthesis of haemoglobin and red blood cells. Vitamin B₁ is an appetite stimulant. In addition, Vitamin B₆ contributes to optimal bone and muscle development. Biotin (B₇) is essential for good skin and hair health. Lastly, Vitamins B₁, B₂, B₃ and B₅ intervene in metabolic processes that transform nutrients into energy
 - **Vitamin C** (Ascorbic Acid) is highly antioxidant, neutralises free radicals and helps prevent aging-related diseases, cellular stress and joint degeneration. It also helps the functioning of the immune system.
 - Proper intake of **Vitamin D₃** is key for the absorption and normal use of calcium and phosphorus, contributes to bone, dental and muscle health, and the optimal functioning of the immune system.
 - **Vitamin E** is a powerful antioxidant, avoids damage to cellular membranes and protects tissues from free radicals. It also enhances the immune system.
- Macro-minerals are those that are needed in greater quantities.
 - **Calcium** and **Phosphorus** intervene in the formation and metabolism of bones and teeth. Calcium, on the other hand, intervenes in many other functions, such as nerve impulse transmission, muscle contraction and coagulation; proper intake is essential in growth phases and in pregnant and lactating females. Phosphorus forms part of the cellular membranes and DNA and RNA molecules.
 - **Potassium** is essential for proper cellular function; it ensures the balance between interior and exterior pressure and intervenes in energetic metabolism. It is also implicated in proper heart function.
 - **Magnesium** is involved in nerve conduction and muscle contractions; its deficit may cause the appearance of neurological disorders.
 - Micro-minerals are those which are necessary in smaller amounts.
 - **Iron** and **Copper** are essential for the synthesis of haemoglobin and red blood cells. They participate in numerous enzymatic processes related to respiration and cellular oxidation. Copper also intervenes in the synthesis of collagen and myelin.
 - **Zinc** is indispensable for the synthesis of collagen and keratin. In the skin, it actively participates in the regeneration of the extra-cellular matrix, healing processes, repair of connective tissue, inflammation and cellular growth. Correct intake maintains the integrity of the skin and quality of the coat.
 - **Selenium** is a powerful antioxidant; as such, it minimises the damage caused by free radicals produced in situations of aging, intense physical activity or inflammatory diseases and stimulates immune system functionality.
 - **Iodine** is involved in the metabolism and synthesis of thyroid hormones.
 - **Omega-3** fatty acids produce many beneficial effects in the body; they possess an anti-inflammatory activity inhibiting the synthesis of certain chemical mediators of inflammation. They are associated with the treatment of chronic inflammatory diseases such as allergies, asthma, arthrosis and chronic kidney disease. In addition, they improve the skin's elasticity.
 - **Linoleic Acid (Omega-6)** is an essential fatty acid important for a multitude of processes. It forms part of the cellular membrane, being indispensable for its synthesis, and its deficit may increase capillary fragility and, as a result, hair loss. Lack of Omega-6 also affects the integrity of the skin barrier, causing poor skin condition and higher sensitivity, increasing the appearance of skin pathologies. It also intervenes in bone and dental development and good joint function.
 - **Taurine** is an essential amino acid in cats, involved in the proper functioning of the cardiac muscle, vision and reproduction. It is also necessary for the synthesis of bile salts by the liver. Its deficit has been associated with severe heart pathologies, blindness and reproductive problems. It also has an antioxidant effect.
 - **Methionine** is also an essential amino acid that must be administered in the diet, since it cannot be synthesised by the body. It is an intermediary in the synthesis of other amino acids like cystine. It plays an important role in hepatic metabolism, protecting the liver from the toxic effects of hepatotoxins. It also has an antioxidant action. It helps to improve hepatic function in senior dogs or those under intense or prolonged drug therapy.
 - **Cystine** is indispensable for the synthesis of keratin; its deficiency produces excessive hair loss and poor coat condition.



MULTIVA® Active Cat

Multi-Vitamin-Mineral Supplement with Amino Acids, Omega-3, Omega-6 and Digestive Enzymes Formulated into Highly Palatable Chews.



Data Sheet

- Digestive enzymes contribute to the digestion, absorption and use of different molecules that the animal ingests. **Bromelain, Protease, Pepsin and Papain** are involved in the digestion of proteins and the assimilation of their amino acids. **Lipase** helps the absorption of fats, enhancing absorption through the intestinal wall. Its administration promotes the improvement of digestive disorders that occur with poor digestion, malabsorption and weight loss.

Indications:

- Improves appetite and general condition in those animals that present apathy, loss of appetite and enervation, while stimulating metabolism.
- Recommended in animals with high nutritional needs due to organic problems (disease, convalescence), stress or high activity.
- Strengthens the immune system, improving defences in situations of stress, environmental changes, convalescence...
- Indicated in older animals to protect and improve health.
- In growing animals to promote optimal bone development and in pregnant and lactating females to improve foetal development and support lactation.
- Helps prevent and fight excessive hair loss and other dermatological problems.
- Recommended in intestinal problems that occur with digestive issues or weight loss.
- Prevents possible vitamin, mineral, amino acid and essential fatty acid deficiencies associated with poor nutrition.

Target Species: Cats.

Safety: MULTIVA® Active Cat is a highly safe product, suitable for all breeds and ages. In cats with special needs it may be necessary to administer for prolonged periods, in which case, follow the instructions of your veterinarian.

Dosage: 1 Chew a day in kittens and small breed adult cats; 2 Chews a day in medium and large breed adult cats. Dosage can be divided and taken twice a day (morning and evening).

Warnings: VetNova is a pioneer in the development of Chews technology to facilitate the administration of supplements to dogs and cats. Unlike tablets, capsules, etc, which are administered orally "by force" to ensure the product is taken, Chews should be freely administered in the feeding bowl and the animal left to consume them voluntarily. Some shy cats may need more time to fully accept them, but once they do, daily intake is easier and more satisfactory. To facilitate initial acceptance, the following strategies can be followed during the first week: 1) Reduce and then progressively increase the dose, 2) Divide the daily dose into two parts (morning and night), 3) Crush the Chew and mix with pâté or any food that appeals to the pet, etc. Keep the container tightly closed, in a cool, dry place, protected from direct sunlight and out of the reach of children and animals.

Presentation: 45 Chews.

Bibliography:

- Alexander JE, Moore MP, Wood LLH. Comparative growth studies in Labrador retrievers fed 5 commercial calorie-dense diets. *Modern Veterinary practice* 1988; 31: 144-148.
- Backus RC. Low plasma taurine concentration in Newfoundland dogs is associated with plasma methionine and cyst(e)ine concentrations and low taurine synthesis. *J. Nutr.* 2006; 136: 2525-2533.
- Backus RC, Cohen G, Pion PD, et al. Taurine deficiency in Newfoundlands fed commercially balanced diets. *Journal of the American Medical Association* 2003; 223 (8): 1130-1136.
- Bai SC, Sampson DA, Morris JG, Rogers QR. The level of dietary protein affects vitamin B-6 requirement of cats. *J. Nutr.* 1991; 121: 1054-1061.
- Bai SC, Sampson DA, Morris JG, Rogers QR. Vitamin B-6 requirement of growing kittens. *J. Nutr.* 1989; 119: 1020-1027 b.



MULTIVA® Active Cat

Multi-Vitamin-Mineral Supplement with Amino Acids, Omega-3, Omega-6 and Digestive Enzymes Formulated into Highly Palatable Chews.



Data Sheet

- Bauer JE. Fatty acid metabolism in domestic cats (*Felis catus*) and cheetahs (*Acinonyx jubatus*). Proceedings of the Nutrition Society 1997; 56: 1013–1024.
- Bauer JE. Responses of dogs to dietary omega-3 fatty acids. J Amer Vet Med Assoc 2007; 11: 1657–1661.
- Bauer J, Heinemann KM, Lees GE, Waldron MK. Retinal functions of young dogs are improved and maternal plasma phospholipids are altered with diets containing long-chain n-3 PUFA during gestation, lactation and a er weaning. J. Nutr. 2006; 136: 1991S-1994S.
- Bauer JE, Heinemann KM, Bigley KE, et al. Maternal diet alphanolenic acid during gestation and lactation does not increase docosahexaenoic acid in canine milk. J. Nutr. 2004; 134 (8S): 2035S-2038S.
- Bauer JE, Heinemann KM, Lees GE, Waldron MK. Docosahexaenoic acid accumulates in plasma of canine puppies raised on α -linolenic acid-rich milk during suckling but not when fed α -linolenic acid-rich diets a er weaning. J. Nutr. 2006; 136: 2087S-2089S.
- Boemke W, Palm U, Kaczmarczyk G, Reinhardt HW. Effect of high sodium and high water intake on 24 h-potassium balance in dogs. Z. Versuchstierkd. 1990; 33 (4): 179-185.
- Booles D, Burger IH, Whyte AL, et al. Effects of two levels of zinc intake on growth and trace element status in Labrador puppies. J Nutr 1991; 121: S79-S80.
- Brown SA, Brown CA, Crowell WA, et al. Effects of dietary polyunsaturated fatty acid supplementation in early renal insufficiency in dogs. J Lab Clin Med 2000;135:275–286.
- Burger I. Water balance in the dog and the cat. Pedigree Digest 1979; 6: 10-11.
- Burger, IH, and Barnett, K. C. The taurine requirement of the adult cat J. Small. Anim. Pract. 1982; 23: 533-537.
- Burger IH, Smith P. Aminosäurenbedarf erwachsener Katzen. In: Ernährung, Fehlernährung, und Diätetik bei Hund und Katze – Proceedings of the International Symposium Hannover (DE), September 3-4, 1987: pp. 93-97.
- Carlomagno G and Unfer V. Inositol safety: clinical evidences. European Review for Medical and Pharmacological Sciences 2011; 15, 931–936.
- Castillo VA, Lalia JC, Junco M, et al. Changes in thyroid function in puppies fed a high iodine commercial diet. Veterinary Journal 2001; 161 (1): 80-84.
- Castillo VA, Pisarev MA, Lalia JC, et al. Commercial diet induced hypothyroidism due to high iodine. A histological and radiological analysis. Veterinary Quarterly 2001; 23 (4): 218-223.
- Cline JL, Czarniecki-Maulden, Losonsky JM, et al. Effect of increasing dietary vitamin A on bone density in adult dogs. J. Anim. Sci. 1997; 75: 2980-2985.
- Cline JL, Odle J, Easter RA. The riboflavin requirement of adult dogs at maintenance is greater than previous estimates J Nutr. 1996 Apr; 126 (4):984-988.
- Czarniecki-Maulden GL, Deming JG, Izquierdo JV. Evaluation of practical dry dog foods suitable for all life stages. J. Amer. Vet. Med. Assoc. 1989; 195 (5): 583-590.
- Deady JE, Anderson B, O'Donnell III JA, et al. Effects of level of dietary glutamic acid and thiamine on food intake, weight gain, plasma amino acids and thiamin status of growing kittens. J. Nutr. 1981; 111: 1568-1579.
- Deady JE, Rogers QR, Morris JG. Effect of high dietary glutamic acid on the excretion of 35S-thiamin in kittens. J. Nutr. 1981; 111: 1580-1585.
- Delaney SJ, Kass PH, Rogers QR and AJ Fascetti. Plasma and whole blood taurine in normal dogs of varying size fed commercially prepared food. Journal of Animal Physiology and Animal Nutrition 2003; 87: 236-344.
- Douglass GM, Fern EB, Brown RC. Feline plasma and whole blood taurine levels as influenced by commercial dry and canned diets. J. Nutr. 1991; 121: 179S-180S.
- Dunbar BL, Bauer JE. Conversion of essential fatty acids by delta 6-desaturase in dog liver microsomes. J Nutr 2002; 132: 1701S–1703S.
- Earle KE, Smith PM. The effect of taurine content on the plasma taurine concentration of the cat Brit. J. Nutr. 1991; 66: 227-235.
- Fascetti AJ, Morris JG, Rogers QR. Dietary copper influences reproductive efficiency of queens. J. Nutr 1998; 128: 2590S-2592S.
- Fascetti AJ, Reed JR, Rogers QR, and Backus RC. Taurine deficiency in dogs with dilated cardiomyopathy: 12 cases (1997-2001). JAVMA 2003; 223 (8): 1137-1141.
- Fox PR. Taurine deficiency dilated cardiomyopathy and idiopathic myocardial failure. In: Textbook of Veterinary Internal Medicine. SJ Ettinger, EC Feldman Edits. 5th edition, WB Saunders Company Philadelphia, PA. 2000: pp. 908-912.
- Freeman LM, Rush JE, Kehayias JJ. Nutritional alterations and the effects of fish oil supplementation in dogs with heart failure. J Vet Intern Med 1998; 12: 440–448.
- Freeman LM, Rush JE, Markwell PJ. Effects of dietary modification in dogs with early chronic valvular disease. J Vet Intern Med 2006; 20: 1116–1126.



VetNova

MULTIVA® Active Cat

Multi-Vitamin-Mineral Supplement with Amino Acids, Omega-3, Omega-6 and Digestive Enzymes Formulated into Highly Palatable Chews.



Data Sheet

- Freytag TL, Liu SM, Rogers AR, Morris JG. Teratogenic effects of chronic ingestion of high levels of vitamin A in cats. *J. Anim Phys and Anim Nutr.* 2003; 87: 42-51.
- Goldy GG, Burr JR, Longardner CN et al. Effects of measured doses of vitamin A fed to healthy dogs for 26 weeks. *Veterinary Clinical Nutrition* 1996; 3 (2): 42-49.
- Goodman SA, Montgomery RD, Fitch RB et al. Serial orthopaedic examinations of growing great Dane puppies fed three diets varying in calcium and phosphorus. In: *Recent advances in canine and feline nutrition. Vol 2. Iams Nutrition Symposium Proceedings.* G. Reinhardt & D. Carye edits. Wilmington, Ohio, Orange Frazer Press. 1998; pp. 3-12.
- Hall JA. Potential adverse effects of long-term consumption of (n3) fatty acids. *Comp Cont Educ Pract Vet.* 1996; 18 (8): 879-895.
- Hall JA, Tooley KA, Gradin JL, et al. Influence of dietary n-6 and n-3 fatty acids and vitamin E on the immune response of healthy geriatric dogs. *Am J Vet Res* 2003; 64 (6): 762-772.
- Hansen RA, Harris ME, Pluhar GE, et al. Fish oil decreases matrix metalloproteinases in knee synovia of dogs with inflammatory joint disease. *J Nutr Biochem* 2008;19(2):101-108.
- Hazewinkel HAW. Influences of different calcium intakes on calcium metabolism and skeletal development in young Great Danes. Thesis Utrecht University, 1985.
- Heinemann KM, Bauer JE. Timely Topics in Nutrition - Docosahexaenoic acid and neurologic development in animals. *J. Am Vet Med Assoc* 2006; 228 (5): 700-705.
- Heinemann KM, Waldron MK, Bigley KE, Bauer JE. Improvement of retinal function in canine puppies from mothers fed dietary long chain n-3 polyunsaturated fatty acids during gestation and lactation. *J Vet Int Med* 2005; 19 (3): 442-443, Abstr. 155.
- Heinemann KM, Waldron MK, Bigley KE, et al. Long-Chain (n-3) Polyunsaturated fatty acids are more efficient than α -linolenic acid in improving electroretinogram responses of puppies exposed during gestation, lactation, and weaning. *J. Nutr.* 2005; 135: 1960-1966.
- Hendriks WH, Wu YB, Shields RG, et al. Vitamin E requirement of adult cats increases slightly with high dietary intake of polyunsaturated fatty acids. *J Nutr* 2002; 132: 1613S-1615S.
- Hickman MA, Rogers QR, Morris JG. Effect of Processing on Fate of Dietary [14C] Taurine in Cats. *J. Nutrition* 1990; 120: 995-1000.
- Hickman MA, Rogers QR, Morris JG. Taurine Balance is Different in Cats Fed Purified and Commercial Diets. *J. Nutr.* 1992; 122: 553-559.
- Kelley R, Lepine A. Improving puppy trainability through nutrition. *Proceedings from Iams Clinical Nutrition Symposium 2005, Seville, Spain.*
- Kronfeld DS, Biotin and Avidin. In *vitamin & Mineral Supplementation for dogs and cats – A monograph on micronutrients* Veterinary Practice Publishing Company 1989: 71-72, 99.
- Laflamme DP. Effect of breed size on calcium requirements for puppies. *Supplement to Compendium on Continuing Education for the Practicing Veterinarian* 2001; 23 (9A): 66-69.
- Lauten SD, Cox NR, Brawner WR, et al. Influence of dietary calcium and phosphorus content in a fixed ration on growth and development of Great Danes. *Am J Vet Res.* 2002; 63 (7): 1036-1047.
- Logas D, Beale KM, Bauer JE. Potential benefits of dietary supplementation with marine-life oil. *J Am Vet Med Assoc* 1991;199(11):1631-1636.
- Morris JG, Rogers QR, Kim SW, Backus RC. Dietary taurine requirement of cats is determined by microbial degradation of taurine in the gut. *Vet. Clin. Nutr.* 1994; 1 (3): 118-127.
- Morris PJ, Salt C, Raila J, et al. Safety evaluation of vitamin A in growing dogs. *Br J Nutr* 2012; 108: 1800-1809.
- NRC (National Research Council). *Nutrient Requirements of Dogs and Cats.* The National Academic Press, Washington, DC. 2006. Washington, DC. 2006.
- Oddle J, Roach M, Baker DH. Taurine utilization by cats. *J. Nutr.* 1993; 123: 1932-1933.
- Pastoor FJ, Van't Klooster AT, Opitz R, et al. Effect of dietary magnesium level on urinary and faecal excretion of calcium, magnesium and phosphorus in adult, ovariectomized cats. *Br J Nutr.* 1995; 74 (1): 77-84.
- Pawlosky RJ, Denkins Y, Ward G, et al. Retinal and brain accretion of long-chain polyunsaturated fatty acids in developing felines: the effects of corn oil-based maternal diets. *Am. J. Clin Nutr* 1997; 65 (2): 465-472.
- Pion PD, Sanderson SL, and Kittleson MD. The effectiveness of taurine and levocarnitine in dogs with heart disease. *Vet Clin of North Am Small Anim Pract* 1998; 1495-1514.
- Pion, Kittleson & Rogers. Myocardial failure in cats associated with low plasma taurine: a reversible cardiomyopathy *Science* 1987; 237: 764-768.
- Re S, Zanoletti M, Emanuele E. Aggressive dogs are characterized by low omega-3 polyunsaturated fatty acid status. *Vet Res Commun* 2008;32:225-230.
- Saker KE, Eddy AL, Thatcher CD, Kalnitsky J. Manipulation of Dietary (n-6) and (n-3) Fatty Acids Alters Platelet



MULTIVA® Active Cat

Multi-Vitamin-Mineral Supplement with Amino Acids, Omega-3, Omega-6 and Digestive Enzymes Formulated into Highly Palatable Chews.



Data Sheet



- Function in Cats. *J Nutr* 1998;128:2645S–2647S.
- Sanderson SL, Gross KL, Ogburn PN, et al. Effects of dietary fat and L-carnitine on plasma and whole-blood taurine concentrations and cardiac function in healthy dogs fed protein-restricted diets. *Am J Vet Res.* 2001; 62: 1616-1623.
 - Schoenmakers I, Hazewinkel HAW, Voorhout G, et al. Effect of diets with different calcium and phosphorus contents on the skeletal development and blood chemistry of growing grate Danes. *Vet Rec.* 2000; 147: 652-660.
 - Schweigert FJ, Bok V. Vitamin A in blood plasma and urine of dogs is affected by the dietary level of vitamin A. *Int J Vitam Nutr Res* 2000; 70, 84-91.
 - Seawright AA, English PB, Gartner RJW. Hypervitaminosis A and deforming cervical spondylosis of the cat. *J. Comp. Path.* 1967; 77: 29-39.
 - Sih TR, Morris JG, Hickman MA. Chronic ingestion of high concentrations of cholecalciferol in cats. *Am. J. Vet. Res.* 2001; 62 (9): 1500-1506.
 - Smith CE, Freeman LM, Rush, JE, et al. Omega-3 fatty acids in Boxer dogs with arrhythmogenic right ventricular cardiomyopathy. *J Vet Intern Med* 2007;21:256–273.
 - Smith GA, Mayhew PD, Kapatkin AS, et al. Reevaluation of risk factors for degenerative joint disease associated with hip dysplasia in German Shepherd Dogs, Golden Retrievers, Labrador Retrievers, and Rottweilers. *J Am Vet Med Assoc* 2001;219:1719–1724.
 - Spitze A.R, Wong D.L, Rogers Q.R, Fascetti A.J. (2003) Taurine concentrations in animal feed ingredients; cooking influences taurine content. *Journal of Animal Physiology and Animal Nutrition* 87: 251-262.
 - Stratton-Phelps M, Backus RC, Rogers QR, and Fascetti AJ. Dietary rice bran decreases plasma and whole-blood taurine in cats. *J. Nutr.* 2002; 132: 1745S-1747S.
 - Strieker MJ, Morris JG, Feldman BF, Rogers QR. Vitamin K deficiency in cats fed commercial fish-based diets. *J Small Anim Pract.* 1996; 37 (7): 322-326.
 - Törres CL, Backus RC, Fascetti AJ, Rogers QR. Taurine status in normal dogs fed a commercial diet associated with taurine deficiency and dilated cardiomyopathy. *Journal of Animal Physiology and Animal Nutrition* 2003; 87: 359-372.
 - Tryfonidou MA, Holl MS, Vastenburg M, et al. Chapter 7. Moderate vitamin D3 supplementation mildly disturbs the endochondral ossification in growing dogs. In: *PhD Thesis Utrecht University* 19 December 2002: pp. 110-122.
 - Tryfonidou MA, Stevenhagen JJ, van den Bemd GJCM, et al. Moderate cholecalciferol supplementation depresses intestinal calcium absorption in growing dogs. *J. Nutr.* 2002; 132: 2644-2650.
 - Wander RC, Hall JA, Gradin JL, et al. The ratio of dietary (n-6) to (n-3) fatty acids influences immune system function, eicosanoid metabolism, lipid peroxidation and vitamin E in aged dogs. *J. Nutr.* 1997; 127: 1198-1997.
 - Watson AL, Fray TR, Bailey J, Baker CB, Beyer SA and Markwell PJ. Dietary constituents are able to play a beneficial role in canine epidermal barrier function. *Experimental Dermatology*; 2006: 15, 74–81.
 - Weber M, Martin L, Dumon H, et al. Growth and skeletal development in two large breeds fed 2 calcium levels. *J. Vet Int. Med* 2000; 14 (May/June): 388 Abstr. 243.
 - Weber M, Martin L, Dumon H, et al. Calcium in growing dogs of large breed: a safety range? *ESVCN Congress Amsterdam, April 2000, Abstr.*
 - Wedekind K, Beyer R, Combs Jr. G. Is selenium addition necessary in pet foods? *FASEB J.* 1998; 12: Abstr. 823.
 - Wedekind KJ, Blumer ME, Huntington CE, et al. The Feline Iodine Requirement is Lower than the 2006 NRC Recommended Allowance. *J Anim Phys and Anim Nutr* 2010; 94 (4): 527–539.
 - Wedekind K., Combs Jr. G. Selenium in pet foods: Is bioavailability an issue? *Compend Cont Educ Pract Vet* 2000; 22 (Suppl.): 17-22.
 - Wedekind K, Yu S, Combs GF. The selenium requirement of the puppy. *J. Anim. Physiol. a. Anim. Nutr.* 2004; 88: 340-347.
 - Yu S, Morris JG. Sodium requirement of adult cats for maintenance based on plasma aldosterone concentration. *J. Nutr.* 1999; 129: 419-423.
 - Yu S, Morris JG. The minimum sodium requirement of growing kittens defined on the basis of plasma aldosterone concentration. *J. Nutr.* 1997; 127: 494-501.
 - Zentek J, Kohn B, Morris P, et al. Effect of dietary vitamin A on plasma levels and urinary excretion of retinol and retinyl esters and clinical parameters in puppy dogs. In: *Proceedings of the 13th Congress of the ESVCN, Oristano, Italy 15-17 October 2009, p. 97.*

If you are interested in any of the articles listed, please do not hesitate to request them through the following contacts: vetnova@vetnova.net, +34 918 440 273, or your VetNova or Distributor Sales Representative.

VetNova

T.: +34 918 440 273 · vetnova@vetnova.net · www.vetnova.net

VN-PUB-0105EN.0616

